



GENEVA | 31 January 2013 -- **Adults** should **consume less than 2,000 mg of sodium**, or **5 grams**

of

salt

, and at least

3,510 mg

of

potassium

per

day

, according to new guidelines issued by the World Health Organization (WHO). A person with either elevated sodium levels and low potassium levels could be at risk of raised

blood

pressure

which increases the

risk

of

heart disease

and

stroke

.

Sodium is found naturally in a variety of foods, including milk and cream (approximately 50 mg of sodium per 100 g) and eggs (approximately 80 mg/100 g). It is also found, in much higher amounts, in processed foods, such as bread (approximately 250 mg/100 g), processed meats like bacon (approximately 1,500 mg/100 g), snack foods such as pretzels, cheese puffs and popcorn (approximately 1,500 mg/100 g), as well as in condiments such as soy sauce (approximately 7,000 mg/100 g), and bouillon or stock cubes (approximately 20,000 mg/100 g).

Potassium-rich foods include: beans and peas (approximately 1,300 mg of potassium per 100 g), nuts (approximately 600 mg/100 g), vegetables such as spinach, cabbage and parsley (approximately 550 mg/100 g) and fruits such as bananas, papayas and dates (approximately 300 mg/100 g). Processing reduces the amount of potassium in many food products.

Currently, most people consume too much sodium and not enough potassium.

“Elevated blood pressure is a major risk for heart disease and stroke – the number one cause of death and disability globally,” says Dr Francesco Branca, Director of WHO’s Department of Nutrition for Health and Development. “These guidelines also make recommendations for children over the age of 2. This is critical because children with elevated blood pressure often become adults with elevated blood pressure.”

WHO issues new guidance on dietary salt and potassium

Écrit par WHO

Jeudi, 31 Janvier 2013 18:02 - Mis à jour Vendredi, 01 Février 2013 10:34

The guidelines are an important tool for public health experts and policymakers as they work in their specific country situations to address noncommunicable diseases such as heart disease, stroke, diabetes, cancer and chronic respiratory diseases. Public health measures to reduce sodium and increase potassium consumption and thereby decrease the population's risk of high blood pressure and heart disease can include food and product labelling, consumer education, updating national dietary guidelines, and negotiating with food manufacturers to reduce the amount of salt in processed foods.

WHO is also updating guidelines on the intake of fats and sugars associated to reduced risk of obesity and noncommunicable diseases.

- *WHO Guidelines: Sodium intake for adults and children* www.who.int/nutrition/publications/guidelines/sodium_intake
- *WHO Guidelines: Potassium intake for adults and children* www.who.int/nutrition/publications/guidelines/potassium_intake