Écrit par WHO Mardi, 11 Octobre 2016 12:00 - Mis à jour Mardi, 11 Octobre 2016 12:01



11 October, 2016 ¦ Geneva: Taxing sugary drinks can lower consumption and reduce obesity, type 2 diabetes and tooth decay, says a new World Health Organization (WHO) report.

Fiscal policies that lead to at least a 20% increase in the retail price of sugary drinks would result in proportional reductions in consumption of such products, according to the report titled *Fiscal policies for Diet and Prevention of Noncommunicable Diseases* (NCDs).

Reduced consumption of sugary drinks means lower intake of "free sugars" and calories overall, improved nutrition and fewer people suffering from overweight, obesity, diabetes and tooth decay.

Free sugars refer to monosaccharides (such as glucose, fructose) and disaccharides (such as sucrose or table sugar) added to foods and drinks by the manufacturer, cook or consumer, and sugars naturally present in honey, syrups, fruit juices and fruit juice concentrates.

Obesity on the rise

"Consumption of free sugars, including products like sugary drinks, is a major factor in the global increase of people suffering from obesity and diabetes," says Dr Douglas Bettcher, Director of WHO's Department for the Prevention of NCDs. "If governments tax products like sugary drinks, they can reduce suffering and save lives. They can also cut healthcare costs and increase revenues to invest in health services."

In 2014, more than 1 in 3 (39%) adults worldwide aged 18 years and older were overweight. Worldwide prevalence of obesity more than doubled between 1980 and 2014, with 11% of men and 15% of women (more than half a billion adults) being classified as obese.

In addition, an estimated 42 million children aged under 5 years were overweight or obese in 2015, an increase of about 11 million during the past 15 years.[1] Almost half (48%) of these children lived in Asia and 25% in Africa.

The number of people living with diabetes has also been rising, from 108 million in 1980 to 422 million in 2014. The disease was directly responsible for 1.5 million deaths in 2012 alone.

Need to reduce sugar intake

"Nutritionally, people don't need any sugar in their diet. WHO recommends that if people do consume free sugars, they keep their intake below 10% of their total energy needs, and reduce it to less than 5% for additional health benefits. This is equivalent to less than a single serving (at least 250 ml) of commonly consumed sugary drinks per day," says Dr Francesco Branca, Director of WHO's Department of Nutrition for Health and Development.

According to the new WHO report, national dietary surveys indicate that drinks and foods high in free sugars can be a major source of unnecessary calories in people's diets, particularly in the case of children, adolescents and young adults.

WHO urges global action to curtail consumption and health impacts of sugary drinks

Écrit par WHO

Mardi, 11 Octobre 2016 12:00 - Mis à jour Mardi, 11 Octobre 2016 12:01

It also points out that some groups, including people living on low incomes, young people and those who frequently consume unhealthy foods and beverages, are most responsive to changes in prices of drinks and foods and, therefore, gain the highest health benefits.

Using fiscal policies to reduce consumption

Fiscal policies should target foods and beverages for which healthier alternatives are available, the report adds.

The report presents outcomes of a mid-2015 meeting of global experts convened by WHO and an investigation of 11 recent systematic reviews of the effectiveness of fiscal policy interventions for improving diets and preventing NCDs and a technical meeting of global experts. Other findings include:

- Subsidies for fresh fruits and vegetables that reduce prices by 10–30% can increase fruit and vegetable consumption;
- Taxation of certain foods and drinks, particularly those high in saturated fats, trans fat, free sugars and/or salt appears promising, with existing evidence clearly showing that increases in the prices of such products reduces their consumption;
- Excise taxes, such as those used on tobacco products, that apply a set (specific) amount of tax on a given quantity or volume of the product, or particular ingredient, are likely to be more effective than sales or other taxes based on a percentage of the retail price;
- Public support for such tax increases could be increased if the revenue they generate is earmarked for efforts to improve health systems, encourage healthier diets and increase physical activity.

A number of countries have taken fiscal measures to protect people from unhealthy products. These include Mexico, which has implemented an excise tax on non-alcoholic beverages with added sugar, and Hungary, which has imposed a tax on packaged products with high sugars, salt or caffeine levels.

Countries, such as the Philippines, South Africa and the United Kingdom have also announced intentions to implement taxes on sugary drinks.